

Medications to stop SEVEN days prior to procedure:

COUMADIN

If you take Coumadin we need blood work prior to procedure

BLOOD THINNERS

Plavix (Clopidogrel)
Aggrenox (Aspirin and Dipyridamole)
Pradaxa (Dabigatran)
Xarelto (Rivaroxaban)
Effient (Prasugrel)
Brilinta (Ticagrelor)
Eliquis (Apixaban)
Trental (Pentoxil)

ASPIRIN PRODUCTS

Anacin
Excedrin
Bayer
Ecotrin
Alka-Seltzer

HEADACHE POWDERS

BC
Stanback
Goody

IBUPROFEN

Advil
Motrin

**NON-STEROIDAL ANTI-INFLAMMATORY
(NSAIDS)**

Cataflam (Diclofenac Potassium)
Voltaren (Diclofenac Sodium)
Lodine (Etodolac)
Indocine (Indomethacin)
Orudis (Ketoprofen)
Toradol (Ketorolac Tromethamine)
Mobic (Meloxicam)
Relafen (Nabumetone)
Aleve (Naproxen/Naprosyn)
Feldene (Piroxicam)
Clinoril (Sulindac/Sundac)
Daypro (Oxaprozin)
Dolobid (Diflunisal)
Amigesic (Salsalate)
Tolectin (Tolmetin)
Vicoprofen (Hydrocodone and Ibuprofen)

ANTIBIOTICS

No infection
No fever
No flu-like symptoms (cough, body aches)